


Penicillin G Potassium for Injection, USP

Rx only

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Penicillin G Potassium for Injection, USP and other antibacterial drugs, Penicillin G Potassium for Injection, USP should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

DESCRIPTION

Penicillin G Potassium for Injection, USP is a natural penicillin. Penicillin G Potassium for Injection, USP is a sterile, pyrogen-free powder for reconstitution. Penicillin G Potassium for Injection, USP is an antibacterial agent intended for parenteral administration. Chemically, Penicillin G Potassium for Injection, USP is monopotassium (2S,5R,6R)-3,3-dimethyl-7-oxo-6-(2-phenylacetamido)-4-thia-1-azabicyclo (3.2.0) heptane-2-carboxylate. It has a molecular weight of 372.48 and the following structural formula:



Penicillin G potassium is a colorless or white crystal, or a white crystalline powder which is odorless, or practically so, and moderately hygroscopic. Penicillin G potassium is very soluble in water. The pH of the reconstituted product is between 6.0-8.5. Penicillin G potassium is buffered with sodium citrate to an optimum pH.

Penicillin G Potassium for Injection, USP is supplied in vials equivalent to 1,000,000 units (1 million units), 5,000,000 units (5 million units), for intramuscular and intravenous use or 20,000,000 units (20 million units), for intravenous use only, of penicillin G as the potassium salt. Each million units contains approximately 6.9 milligrams of sodium (0.3 mEq) and 65.8 milligrams of potassium (1.68 mEq).

CLINICAL PHARMACOLOGY

After an intravenous infusion of penicillin G, peak serum concentrations are attained immediately after completion of the infusion. In a study of ten patients administered a single 5 million unit dose of penicillin G intravenously over 3 to 5 minutes, the mean serum concentrations were 400 mcg/mL, 273 mcg/mL and 3 mcg/mL at 5 to 6 minutes, 10 minutes and 4 hours after completion of the injection, respectively. In a separate study, five healthy adults were administered one million units of penicillin G intravenously, either as a bolus over 4 minutes or as an infusion over 60 minutes. The mean serum concentration eight minutes after completion of the bolus was 45 mcg/mL and eight minutes after completion of the infusion was 14.4 mcg/mL. The mean β -phase serum half-life of penicillin G administered by the intravenous route in ten patients with normal renal function was 42 minutes, with a range of 31 to 50 minutes.

The clearance of penicillin G in normal individuals is predominantly via the kidney. The renal clearance, which is extremely rapid, is the result of glomerular filtration and active tubular transport, with the latter route predominating. Urinary recovery is reported to be 58 to 85% of the administered dose. Renal clearance of penicillin is delayed in premature infants, neonates and in the elderly due to decreased renal function. The serum half-life of penicillin G correlates inversely with age and clearance of creatinine and ranges from 3.2 hours in infants 0 to 6 days of age to 1.4 hours in infants 14 days of age or older.

Nonrenal clearance includes hepatic metabolism and, to a lesser extent, biliary excretion. The latter routes become more important with renal impairment.

Probenecid blocks the renal tubular secretion of penicillin. Therefore, the concurrent administration of probenecid prolongs the elimination of penicillin G and, consequently, increases the serum concentrations.

Penicillin G is distributed to most areas of the body including lung, liver, kidney, muscle, bone and placenta. In the presence of inflammation, levels of penicillin in abscesses, middle ear, pleural, peritoneal and synovial fluids are sufficient to inhibit most susceptible bacteria. Penetration into the eye, brain, cerebrospinal fluid (CSF) or prostate is poor in the absence of inflammation. With inflamed meninges, the penetration of penicillin G into the CSF improves, such that the CSF/serum ratio is 2 to 6%. Inflammation also enhances its penetration into the pericardial fluid. Penicillin G is actively secreted into the bile resulting in levels at least 10 times those achieved simultaneously in serum. Penicillin G penetrates poorly into human polymorphonuclear leukocytes.

In the presence of impaired renal function, the β -phase serum half-life of penicillin G is prolonged. β -phase serum half-lives of one to two hours were observed in azotemic patients with serum creatinine clearance concentrations <3 mg/100 mL and ranged as high as 20 hours in anuric patients. A linear relationship, including the lowest range of renal function, is found between the serum elimination rate constant and renal function as measured by creatinine clearance.

In patients with altered renal function, the presence of hepatic insufficiency further alters the elimination of penicillin G. In one study, the serum half-lives in two anuric patients (excreting <400 mL urine /day) were 7.2 and 10.1 hours. A totally anuric patient with terminal hepatic cirrhosis had a penicillin half-life of 30.5 hours, while another patient with anemia and liver disease had a serum half-life of 16.4 hours. The dose of the penicillin G should be reduced in patients with severe renal impairment, with additional modifications when hepatic disease accompanies the renal impairment. Hemodialysis has been shown to reduce penicillin G serum levels.

Microbiology

Penicillin G is bactericidal against penicillin-susceptible microorganisms during the stage of active multiplication. It acts by inhibiting biosynthesis of cell-wall mucopeptide. It is not active against the penicillinase-producing bacteria, which include many strains of staphylococci. Penicillin G is highly active *in vitro* against staphylococci (except penicillinase-producing strains), streptococci (groups A, B, C, G, H, L, and M), pneumococci and *Neisseria meningitidis*.

Other organisms susceptible *in vitro* to penicillin G are *Haemophilus gonorrhoeae*, *Corynebacterium diphtheriae*, *Bacillus anthracis*, clostridia, *Actinomyces* species, *Spirillum minus*, *Streptobacillus moniliformis*, *Listeria monocytogenes*, and leptospira; *Treponema pallidum* is extremely susceptible. Some species of gram-negative bacilli were previously considered susceptible to very high intravenous doses of penicillin G (up to 80 million units/day) including some strains of *Escherichia coli*, *Proteus mirabilis*, salmonella, shigella, *Enterobacter aerogenes* (formerly *Aerobacter aerogenes*) and *Alcaligenes faecalis*. Penicillin G is no longer considered a drug of choice for infections caused by these organisms.

Susceptibility Test Methods

When available, the clinical microbiology laboratory should provide the results of *in vitro* susceptibility test results for antimicrobial drugs used in local hospitals and practice areas to the physician as periodic reports that describe the susceptibility profile of nosocomial and community-acquired pathogens. These reports should aid the physician in selecting the most effective antimicrobial.

Dilution Techniques

Quantitative methods are used to determine antimicrobial minimum inhibitory concentrations (MICs). These MICs provide estimates of the susceptibility of bacteria to antimicrobial compounds. The MICs should be determined using a standardized procedure. Standardized procedures are based on dilution method¹ (broth, agar or microdilution) or equivalent using standardized inoculum and concentrations of penicillin. The MIC values should be interpreted according to the criteria in Table 1.

Diffusion Techniques

Quantitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. One such standardized procedure^{2,3} requires the use of standardized inoculum concentrations. This procedure uses paper disks impregnated with 10 units of penicillin to test the susceptibility of microorganisms to penicillin. Interpretation involves correlation of the diameter obtained in the disk test with the MIC for penicillin. Reports from the laboratory providing results of the standard single-disk susceptibility test with a 10 unit penicillin disk should be interpreted according to the following criteria in Table 1.

Table 1: Susceptibility Test Interpretive Criteria for Penicillin^{2,4}

Pathogen	MIC (mcg/mL)			Disk Diffusion (zone diameter in mm) ⁵		
	Susceptible (S) ⁶	Intermediate (I)	Resistant (R)	Susceptible (S) ⁶	Intermediate (I)	Resistant (R)
Staphylococci	<0.12 ^a	-	≥0.25	≥29 ^b	-	<28
<i>Neisseria gonorrhoeae</i> ^c	≤0.06	0.12 - 1	≥2	≥47	27 - 46	≤26
<i>Streptococcus pneumoniae</i> (meningitis)	≤0.06	-	≥0.12	-	-	-
<i>Streptococcus pneumoniae</i> (pneumonia)	≤2	4	≥8	-	-	-
Beta-hemolytic streptococci ^d	≤0.12 ^a	-	-	≥24 ^a	-	-
<i>Streptococcus</i> spp. Viridans group	≤0.12	0.25 - 2	≥4	-	-	-
<i>Listeria monocytogenes</i> ^e	≤2 ^a	-	-	-	-	-
<i>Bacillus anthracis</i> ^f	≤0.12	-	>0.25	-	-	-

^a Organisms for which no values for disk susceptibility appear cannot be reliably tested with this method.

^b Penicillin-resistant strains of staphylococci produce β -lactamase. An induced β -lactamase test should be performed on all *S. aureus* isolates for which the penicillin MIC is \leq 0.12 mcg/mL or zone diameter is \geq 29 mm before reporting as penicillin susceptible. Rare isolates of staphylococci that contain genes for β -lactamase production may not produce a positive induced β -lactamase test. For serious infections requiring penicillin therapy, laboratories should perform MIC tests and induced β -lactamase testing on all subsequent isolates from the same patient.²

^c A positive *N. gonorrhoeae* β -lactamase test predicts one form of resistance to penicillin. Strains with chromosomally mediated resistance can be detected only by agar dilution or disk diffusion susceptibility test methods.² Isolates with zone diameters \leq 19 mm generally produce β -lactamase.^{2,3}

^d Susceptibility testing of penicillins for treatment of β -hemolytic streptococcal infections need not be performed routinely, because non-susceptible isolates are extremely rare in any β -hemolytic streptococcus and have not been reported from *Streptococcus pyogenes*. Any β -hemolytic streptococcal isolate found to be non-susceptible to penicillin should be re-identified, retested, and, if confirmed, submitted to a public health authority.^{2,3}

^e The current absence of resistant isolates precludes defining results other than "susceptible". Isolates yielding results suggestive of "nonsusceptible" should be submitted to a reference laboratory for further testing.

^f *B. anthracis* strains may contain inducible β -lactamases. *In vitro* penicillinase induction studies suggest that penicillin MICs may increase during therapy. However, β -lactamase testing of clinical isolates of *B. anthracis* is unreliable and should not be performed.⁴

Quality Control

Standardized susceptibility test procedures require the use of laboratory control microorganisms to monitor and ensure the accuracy and precision of the supplies and reagents used in the assay, and the techniques of the individuals performing the test. Standard penicillin powder should provide MIC values provided below. For the diffusion technique, the 10 unit penicillin disk should provide the following zone diameters with the quality control strains:

Table 2. *In vitro* Susceptibility Test Quality Control Ranges for Penicillin

Organism (ATCC [®] #)	MIC range mcg/mL	Disk Diffusion range (mm)
<i>Staphylococcus aureus</i> (29213)	0.25 - 2	Not applicable
<i>Staphylococcus aureus</i> (25923)	Not applicable	26 - 37
<i>Streptococcus pneumoniae</i> (49619)	0.25 - 1	24 - 30
<i>Neisseria gonorrhoeae</i> (49226)	0.25 - 1 ^a	26 - 34

^a Using agar dilution method only. No criteria for broth microdilution are available.²

INDICATIONS AND USAGE

Therapy: Penicillin G Potassium for Injection, USP is indicated in the treatment of serious infections caused by susceptible strains of the designated microorganisms in the conditions listed below. Appropriate culture and susceptibility tests should be done before treatment in order to isolate and identify organisms causing infection and to determine their susceptibility to penicillin G.

Therapy with Penicillin G Potassium for Injection, USP may be initiated before results of such tests are known when there is reason to believe the infection may involve any of the organisms listed below; however, once these results become available, appropriate therapy should be continued.

Clinical Indication	Infecting Organism
Septicemia, empyema, pneumonia, pericarditis, endocarditis, meningitis	<i>Streptococcus pyogenes</i> (group A β -hemolytic streptococci), other β -hemolytic streptococci including groups C,H,G,L and M, <i>Streptococcus pneumoniae</i> and <i>Streptococcus pneumoniae</i> species (non-penicillinase producing strains)
Anthrax	<i>Bacillus anthracis</i>
Actinomycosis (cervico-facial disease and thoracic and abdominal disease)	<i>Actinomyces israelii</i>
Botulism (adjunctive therapy to antitoxin), gas gangrene, and tetanus (adjunctive therapy to human tetanus immune globulin)	<i>Clostridium species</i>
Diphtheria (adjunctive therapy to antitoxin and prevention of the carrier state)	<i>Corynebacterium diphtheriae</i>
Erysipelothrix endocarditis	<i>Erysipelothrix rhusiopathiae</i>
Fusospirochetosis (severe infections of the oropharynx [Vincent's], lower respiratory tract and genital area)	<i>Fusobacterium species</i> and spirochetes
Listeria infections including meningitis and endocarditis	<i>Listeria monocytogenes</i>
Pasteurella infections including bacteremia and meningitis	<i>Pasteurella multocida</i>
Haverhill fever	<i>Streptobacillus moniliformis</i>
Rat-bite fever	<i>Spirillum minus</i> or <i>Streptobacillus moniliformis</i>
Disseminated gonococcal infections	<i>Neisseria gonorrhoeae</i> (penicillin-susceptible)
Syphilis (congenital and neurosyphilis)	<i>Treponema pallidum</i>
Meningococcal meningitis and / or septicemia	<i>Neisseria meningitidis</i>
Gram-negative bacillary infections (bacteremias)	Gram-negative bacillary organisms (i.e. <i>Enterobacteriaceae</i>)

Penicillin G is not the drug of choice in the treatment of gram-negative bacillary infections.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Penicillin G Potassium for Injection, USP and other antibacterial drugs, Penicillin G Potassium for Injection, USP should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

CONTRAINDICATIONS

A history of a hypersensitivity (anaphylactic) reaction to any penicillin is a contraindication.

WARNINGS

SEVERE AND OCCASIONALLY FATAL HYPERSENSITIVITY (anaphylactic) REACTIONS HAVE BEEN REPORTED IN PATIENTS ON PENICILLIN THERAPY. THESE REACTIONS ARE MORE LIKELY TO OCCUR IN INDIVIDUALS WITH A HISTORY OF PENICILLIN HYPERSENSITIVITY AND/OR A HISTORY OF SENSITIVITY TO MULTIPLE ALLERGENS. THERE HAVE BEEN REPORTS OF INDIVIDUALS WITH A HISTORY OF PENICILLIN HYPERSENSITIVITY WHO HAVE EXPERIENCED SEVERE REACTIONS WHEN TREATED WITH CEPHALOSPORINS. BEFORE INITIATING THERAPY WITH PENICILLIN G, CAREFUL INQUIRY SHOULD BE MADE CONCERNING PREVIOUS HYPERSENSITIVITY REACTIONS TO PENICILLINS, CEPHALOSPORINS, OR OTHER ALLERGENS. IF AN ALLERGIC REACTION OCCURS, PENICILLIN G SHOULD BE DISCONTINUED AND APPROPRIATE THERAPY INSTITUTED. **SERIOUS ANAPHYLACTIC REACTIONS REQUIRE IMMEDIATE EMERGENCY TREATMENT WITH EPINEPHRINE, OXYGEN, INTRAVENOUS STEROIDS, AND AIRWAY MANAGEMENT, INCLUDING INTUBATION, SHOULD ALSO BE ADMINISTERED AS INDICATED.**

Clostridium difficile associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including Penicillin G Potassium for Injection, USP, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

PRECAUTIONS

General

Penicillin should be used with caution in individuals with histories of significant allergies and/or asthma (see **WARNINGS**). Whenever allergic reactions occur, penicillin should be withdrawn unless, in the opinion of the physician, the condition being treated is life-threatening and amenable only to penicillin therapy. Penicillin G Potassium for Injection, USP by the intravenous route in high doses (above 10 million units) should be administered slowly because of the potential adverse effects of electrolyte imbalance from the potassium content of the penicillin. Penicillin G Potassium for Injection, USP contains approximately 6.9 milligrams of sodium (0.3 mEq) and 65.8 milligrams of potassium (1.68 mEq) per million units of penicillin G. The use of antibiotics may promote overgrowth of nonsusceptible organisms, including fungi. Inwelling intravenous catheters encourage superinfections. Serious superinfection occur, appropriate measures should be taken. When indicated, incision and drainage or other surgical procedures should be performed in conjunction with antibiotic therapy.

Prescribing Penicillin G Potassium for Injection, USP in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

Information for Patients

Patients should be counseled that antibacterial drugs including Penicillin G Potassium for Injection, USP should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Penicillin G Potassium for Injection, USP is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better after a few days in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may: (1) decrease the effectiveness of the immediate therapy and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Penicillin G Potassium for Injection, USP or other antibacterial drugs in the future.

Diarrhea is a common problem caused by antibiotics which usually ends when the antibiotic is discontinued. Sometimes after starting treatment with antibiotics, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibiotic. If this occurs, patients should contact their physician as soon as possible.

Laboratory Tests

Periodic assessment of organ system function, including frequent evaluation of electrolyte balance, hepatic, renal and hematopoietic systems, and cardiac and vascular status should be performed during prolonged therapy with high doses of intravenous penicillin G (see **ADVERSE REACTIONS**). If any impairment of function is suspected or known to exist, a reduction in the total dosage should be considered (see **DOSAGE AND ADMINISTRATION**). In suspected staphylococcal infections, proper laboratory studies, including susceptibility tests should be performed. All infections due to Group A beta-hemolytic streptococci should be treated for at least 10 days.

Patients being treated for gonococcal infection should have a serologic test for syphilis before receiving penicillin. All cases of penicillin treated syphilis should receive adequate follow-up including clinical and serological examinations. The recommended follow-up varies with the stage of syphilis being treated.

Drug Interactions

Bacteriostatic antibacterials (i.e., chloramphenicol, erythromycin, sulfonamides or tetracyclines) may antagonize the bactericidal effect of penicillin, and concurrent use of these drugs should be avoided. This has been documented *in vivo*, however, the clinical significance of this interaction is not well-documented.

Penicillin blood levels may be prolonged by concurrent administration of probenecid which blocks the renal tubular secretion of penicillins. Other drugs may compete with penicillin G for renal tubular secretion and thus prolong the serum half-life of penicillin. These drugs include: aspirin, phenylbutazone, sulfonamides, indomethacin, thiazide diuretics, furosemide and ethacrynic acid.

Drug/Laboratory Test Interactions

After treatment with penicillin G, a false-positive reaction for glucose in the urine may occur with Benedict's solution, Fehling's solution or Clinites[®] tablet, but not with the enzyme-based tests, such as Clinistix[®] and Tes-Tape[®].

Penicillin G has been associated with pseudoproteinuria by certain test methods.

Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term animal studies have been conducted with this drug.

Pregnancy

Teratogenic Effects

Pregnancy Category B

Penicillin G Potassium for Injection, USP

Rx only

Reproduction studies performed in the mouse, rat, and rabbit have revealed no evidence of impaired fertility or harm to the fetus due to penicillin G. Human experience with the penicillins during pregnancy has not shown any positive evidence of adverse effects on the fetus. There are, however, no adequate and well-controlled studies in pregnant women showing conclusively that harmful effects of these drugs on the fetus can be excluded. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

Penicillins are excreted in human milk. Caution should be exercised when penicillins are administered to a nursing woman.

Pediatric Use

Incompletely developed renal function in newborns may delay elimination of penicillin; therefore, appropriate reductions in the dosage and frequency of administration should be made in these patients. All newborns treated with penicillins should be monitored closely for clinical and laboratory evidence of toxic or adverse effects (see **PRECAUTIONS**).

Pediatric doses are generally determined on a weight basis and should be calculated for each patient individually. Recommended guidelines for pediatric dosages are presented in **DOSAGE AND ADMINISTRATION**.

Geriatric Use

Clinical studies of penicillin G potassium did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

Penicillin G potassium contains 6.9 milligrams of sodium (0.3 mEq) and 65.8 milligrams of potassium (1.68 mEq) per million units of penicillin G. At the usual recommended doses, patients would receive between 6.9 and 165.6 mg/day (0.3 and 7.2 mEq) of sodium. The geriatric population may respond with a blunted natriuresis to salt loading. This may be clinically important with regard to such diseases as congestive heart failure.

ADVERSE REACTIONS

Body As a Whole

The Jarisch-Herxheimer reaction is a systemic reaction, that may occur after the initiation of penicillin therapy in patients with syphilis or other spirochetal infections (i.e., Lyme disease and Relapsing fever). The reaction begins one to two hours after initiation of therapy and disappears within 12 to 24 hours. It is characterized by fever, chills, myalgias, headache, exacerbation of cutaneous lesions, tachycardia, hyperventilation, vasodilation with flushing and mild hypotension. The pathogenesis of the Herxheimer reaction may be due to the release from the spirochetes of heat-stable pyrogen.

Hypersensitivity Reactions

The reported incidence of allergic reactions to all penicillins ranges from 0.7 to 10 percent in different studies (see **WARNINGS**). Sensitization is usually the result of previous treatment with a penicillin, but some individuals have had immediate reactions when first treated. In such cases, it is postulated that prior exposure to penicillin may have occurred via trace amounts present in milk or vaccines.

Two types of allergic reactions to penicillin are noted clinically – immediate and delayed. Immediate reactions usually occur within 20 minutes of administration and range in severity from urticaria and pruritus to angioneurotic edema, laryngospasm, bronchospasm, hypotension, vascular collapse and death (see **WARNINGS**). Such immediate anaphylactic reactions are very rare and usually occur after parenteral therapy, but a few cases of anaphylaxis have been reported following oral therapy. Another type of immediate reaction, an accelerated reaction, may occur between 20 minutes and 48 hours after administration and may include urticaria, pruritus, fever and, occasionally, laryngeal edema.

Delayed reactions to penicillin therapy usually occur within 1 to 2 weeks after initiation of therapy. Manifestations include serum sickness-like symptoms, i.e., fever, malaise, urticaria, myalgia, arthralgia, abdominal pain and various skin rashes, ranging from maculopapular eruptions to exfoliative dermatitis.

Contact dermatitis has been observed in individuals who prepare penicillin solutions.

Gastrointestinal System

Pseudomembranous colitis has been reported with the onset occurring during or after penicillin G treatment. Nausea, vomiting, stomatitis, black or hairy tongue, and other symptoms of gastrointestinal irritation may occur, especially during oral therapy.

Hematologic System

Reactions include neutropenia, which resolves after penicillin therapy is discontinued; Coombs-positive hemolytic anemia, an uncommon reaction, occurs in patients treated with intravenous penicillin G in doses greater than 10 million units/day and who have previously received large doses of the drug; and with large doses of penicillin, a bleeding diathesis, can occur secondary to platelet dysfunction.

Metabolic

Penicillin G potassium (1 million units contains 0.3 mEq of sodium and 1.68 mEq of potassium) may cause serious and even fatal electrolyte disturbances, i.e., hyperkalemia, when given intravenously in large doses.

Neurologic System

Neurotoxic reactions including hyperreflexia, myoclonic twitches, seizures and coma have been reported following the administration of massive intravenous doses, and are more likely in patients with impaired renal function.

Urogenital System

Renal tubular damage and interstitial nephritis have been associated with large intravenous doses of penicillin G. Manifestations of this reaction may include fever, rash, eosinophilia, proteinuria, eosinophiluria, hematuria and in a serum urea nitrogen. Discontinuation of penicillin G results in resolution in the majority of patients.

Local Reactions

Phlebitis and thrombophlebitis may occur, and pain at the injection site has been reported with intravenous administration.

To report SUSPECTED ADVERSE EVENTS, contact **WG Critical Care, LLC at 1-866-562-4708 or the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch**.

OVERDOSAGE

Dose related toxicity may arise with the use of massive doses of intravenous penicillins (40 to 100 million units per day), particularly in patients with severe renal impairment (see **PRECAUTIONS**). The manifestations may include agitation, confusion, asterix, hallucinations, stupor, coma, multifocal myoclonus, seizures and encephalopathy. Hyperkalemia is also possible (see **ADVERSE REACTIONS, Metabolic**).

In case of overdosage, discontinue penicillin, treat symptomatically and institute supportive measures as required. If necessary, hemodialysis may be used to reduce blood levels of penicillin G, although the degree of effectiveness of this procedure is questionable.

DOSAGE AND ADMINISTRATION

Penicillin G potassium may be given intravenously or intramuscularly. The usual dose recommendations are as follows:

Adult Patients

Clinical Indication	Dosage
Serious infections due to susceptible strains of streptococci (including <i>S. pneumoniae</i>) -septicemia, empyema, pneumonia, pericarditis, endocarditis and meningitis	12 to 24 million units/day depending on the infection and its severity administered in equally divided doses every 4 to 6 hours.
Serious infections due to susceptible strains of staphylococci -septicemia, empyema, pneumonia, pericarditis, endocarditis and meningitis	5 to 24 million units/day depending on the infection and its severity administered in equally divided doses every 4 to 6 hours.
Anthrax	Minimum of 8 million units/day in divided doses every 6 hours. Higher doses may be required depending on susceptibility of organism.
Actinomycosis	
Cervico-facial disease	1 to 6 million units/day (*)
Thoracic and abdominal disease	10 to 20 million units/day (*)
Clostridial Infections	
Botulism (adjunctive therapy to antitoxin)	20 million units/day (*)
Gas gangrene (debridement and/or surgery as indicated)	
Tetanus (adjunctive therapy to human tetanus immune globulin)	
Diphtheria (adjunctive therapy to antitoxin and for the prevention of the carrier state)	2 to 3 million units/day in divided doses for 10 to 12 days (*)
Erysipelothrix endocarditis	12 to 20 million units/day for 4 to 6 weeks (*)
Fusospirochetosis (severe infections of the oropharynx [Vincent's], lower respiratory tract and genital area)	5 to 10 million units/day (*)
Listeria infections	
Meningitis	15 to 20 million units/day for 2 weeks (*)
Endocarditis	15 to 20 million units